

HAIR ELEMENTS



LAB#: H061205-0496-1
 PATIENT: Katherine Spehar
 SEX: Female
 AGE: 36

CLIENT#: 24237
 DOCTOR: Justin Caulfield, MD
 Direct Laboratory Services
 300 Mariners Plaza #320
 Mandeville, LA 70448

| POTENTIALLY TOXIC ELEMENTS | | | | |
|----------------------------|------------------------|-----------------|------------------|------------------|
| TOXIC ELEMENTS | RESULT $\mu\text{g/g}$ | REFERENCE RANGE | PERCENTILE | |
| | | | 68 th | 95 th |
| Aluminum | 2.9 | < 7.0 | | |
| Antimony | < 0.01 | < 0.050 | | |
| Arsenic | 0.066 | < 0.060 | | |
| Beryllium | < 0.01 | < 0.020 | | |
| Bismuth | 0.17 | < 0.10 | | |
| Cadmium | 0.035 | < 0.10 | | |
| Lead | 0.79 | < 1.0 | | |
| Mercury | 2.4 | < 1.1 | | |
| Platinum | < 0.003 | < 0.005 | | |
| Thallium | < 0.001 | < 0.010 | | |
| Thorium | < 0.001 | < 0.005 | | |
| Uranium | 0.025 | < 0.060 | | |
| Nickel | 0.09 | < 0.40 | | |
| Silver | 0.05 | < 0.15 | | |
| Tin | 0.04 | < 0.30 | | |
| Titanium | 0.37 | < 1.0 | | |
| Total Toxic Representation | | | | |

| ESSENTIAL AND OTHER ELEMENTS | | | | | | | |
|------------------------------|------------------------|-----------------|-------------------|------------------|------------------|------------------|--------------------|
| ELEMENTS | RESULT $\mu\text{g/g}$ | REFERENCE RANGE | PERCENTILE | | | | |
| | | | 2.5 th | 16 th | 50 th | 84 th | 97.5 th |
| Calcium | 458 | 300- 1200 | | | | | |
| Magnesium | 37 | 35- 120 | | | | | |
| Sodium | 92 | 12- 90 | | | | | |
| Potassium | 23 | 8- 38 | | | | | |
| Copper | 17 | 12- 35 | | | | | |
| Zinc | 250 | 140- 220 | | | | | |
| Manganese | 0.09 | 0.15- 0.65 | | | | | |
| Chromium | 0.31 | 0.20- 0.40 | | | | | |
| Vanadium | 0.022 | 0.018- 0.065 | | | | | |
| Molybdenum | 0.042 | 0.028- 0.056 | | | | | |
| Boron | 0.13 | 0.30- 2.0 | | | | | |
| Iodine | 0.16 | 0.25- 1.3 | | | | | |
| Lithium | < 0.004 | 0.007- 0.023 | | | | | |
| Phosphorus | 161 | 160- 250 | | | | | |
| Selenium | 0.77 | 0.95- 1.7 | | | | | |
| Strontium | 0.47 | 0.50- 7.6 | | | | | |
| Sulfur | 50900 | 44500- 52000 | | | | | |
| Barium | 0.23 | 0.26- 3.0 | | | | | |
| Cobalt | 0.005 | 0.013- 0.050 | | | | | |
| Iron | 7.2 | 5.4- 14 | | | | | |
| Germanium | 0.029 | 0.045- 0.065 | | | | | |
| Rubidium | 0.022 | 0.007- 0.096 | | | | | |
| Zirconium | < 0.007 | 0.020- 0.42 | | | | | |

| SPECIMEN DATA | | | | RATIOS | | |
|------------------|------------|--------------|---------|-----------------|---------------|-----------------------|
| COMMENTS: | | | | ELEMENTS | RATIOS | EXPECTED RANGE |
| Date Collected: | 11/23/2006 | Sample Size: | 0.202 g | Ca/Mg | 12.4 | 4- 30 |
| Date Received: | 12/5/2006 | Sample Type: | Head | Ca/P | 2.84 | 1- 12 |
| Date Completed: | 12/9/2006 | Hair Color: | Brown | Na/K | 4 | 0.5- 10 |
| | | Treatment: | | Zn/Cu | 14.7 | 4- 20 |
| Methodology: | ICP-MS | Shampoo: | | Zn/Cd | > 999 | > 800 |

V06.99

HAIR ELEMENTS REPORT INTRODUCTION

Hair is an excretory tissue for essential, nonessential and potentially toxic elements. In general, the amount of an element that is irreversibly incorporated into growing hair is proportional to the level of the element in other body tissues. Therefore, hair elements analysis provides an indirect screening test for physiological excess, deficiency or maldistribution of elements in the body. Clinical research indicates that hair levels of specific elements, particularly potentially toxic elements such as cadmium, mercury, lead and arsenic, are highly correlated with pathological disorders. For such elements, levels in hair may be more indicative of body stores than the levels in blood and urine.

All screening tests have limitations that must be taken into consideration. The correlation between hair element levels and physiological disorders is determined by numerous factors. Individual variability and compensatory mechanisms are major factors that affect the relationship between the distribution of elements in hair and symptoms and pathological conditions. It is also very important to keep in mind that scalp hair is vulnerable to external contamination of elements by exposure to hair treatments and products. Likewise, some hair treatments (e.g. permanent solutions, dyes, and bleach) can strip hair of endogenously acquired elements and result in false low values. Careful consideration of the limitations must be made in the interpretation of results of hair analysis. The data provided should be considered in conjunction with symptomology, diet analysis, occupation and lifestyle, physical examination and the results of other analytical laboratory tests.

Caution: The contents of this report are not intended to be diagnostic and the physician using this information is cautioned against treatment based solely on the results of this screening test. For example, copper supplementation based upon a result of low hair copper is contraindicated in patients afflicted with Wilson's Disease.

Arsenic High

In general, hair provides a rough estimate of exposure to Arsenic (As) absorbed from food and water. However, hair can be contaminated externally with As from air, water, dust, shampoos and soap. Inorganic As, and some organic As compounds, can cause toxicity. Some research suggests that As may be essential at extremely low levels but its function is not understood. Inorganic As accumulates in hair, nails, skin, thyroid gland, bone and the gastrointestinal tract. Organic As is rapidly excreted in the urine.

As can cause malaise, muscle weakness, vomiting, diarrhea, dermatitis, and skin cancer. Long-term exposure may affect the peripheral nervous, cardiovascular and hematopoietic systems. As is a major biological antagonist to selenium.

Common sources of As are insecticides (calcium and lead arsenate), well water, smog, shellfish (arsenobetaine), and industrial exposure, particularly in the manufacture of electronic components (gallium arsenide).

As burden can be confirmed by urine elements analysis. Comparison of urine As levels pre and post provocation (DMPS, DMSA, D-penicillamine) permit differentiation between recent uptake and body stores.

Bismuth High

No published studies correlate Bismuth (Bi) exposure with hair Bi levels, therefore, hair Bi levels are measured primarily for investigational purposes. Bi is a non-essential element of low toxicity. However, excessive intake of insoluble, inorganic Bi containing compounds can cause nephrotoxicity and encephalopathy. Absorption is dependent upon solubility of the Bi compound, with insoluble Bi excreted in the feces while soluble forms are excreted in the urine. Sources of Bi include: cosmetics (lipstick), Bi containing medications such as ranitidine Bi-citrate, antacids (Pepto Bismol), pigments used in colored glass and ceramics, dental cement, and dry cell battery electrodes.

Symptoms of moderate Bi toxicity include: constipation or bowel irregularity, foul breath, blue/black gum line, and malaise. High levels of Bi accumulation can result in nephrotoxicity (nephrosis, proteinuria) and neurotoxicity (tremor, memory loss, monoclonic jerks, dysarthria, dementia).

Urine elements analysis can be used to corroborate Bi absorption for a period of days or a few weeks after the exposure. Dithiol chelating/complexing agents (DMPS, DMSA) markedly reduced Bi levels in liver and kidneys, and increased Bi in urine in animal studies (J. Lab. Clin. Med.; 119:529-537,1992).

Mercury High

Mercury (Hg) is toxic to humans and animals. The accumulation of Hg in the body is generally reflected by the hair Hg levels, but hair Hg levels can be artifactually high in association with the use of certain hair dyes. Individuals vary greatly in sensitivity and tolerance to Hg burden.

At hair levels below 3 µg/g, Hg can suppress biological selenium function and may cause or contribute to immune dysregulation in sensitive individuals. Hallmark symptoms of excess Hg include: loss of appetite, decreased senses of touch, hearing, and vision, fatigue, depression, emotional instability, peripheral numbness and tremors, poor memory and cognitive dysfunction, and neuromuscular disorders. Hair Hg has been reported to correlate with acute myocardial infarction and on average each 1 µg/g of hair Hg was found to correlate with a 9% increase in AMI risk (Circulation 1995; 91:645-655).

Sources of Hg include dental amalgams, contaminated seafood, water supplies, some hemorrhoidal preparations, skin lightening agents, instruments (thermometers, electrodes, batteries), and combustion of fossil fuels, some fertilizers, and the paper/pulp and gold industries. After dental amalgams are installed or removed a transient (several months) increase in hair Hg is observed. Also, "baseline" hair Hg levels for individuals with dental amalgams are higher (about 1 to 2 µg/g) than are baseline levels for those without (below 1 µg/g).

Confirmatory tests for elevated Hg are measurement of whole blood as an indication of recent/ongoing exposure (does not correlate with whole body accumulation) and measurement of urine Hg following use of a dithiol chelating or mobilizing agent such as DMSA or DMPS (an indication of total body burden).

Sodium High

Sodium (Na) is an essential element with extracellular electrolyte functions. However, these functions do not occur in hair. Hair Na measurement should be considered a screening test only;

blood testing for Na and electrolyte levels is much more diagnostic and indicative of status. High hair Na may have no clinical significance or it may be the result of an electrolyte imbalance. A possible imbalance for which high hair Na is a consistent finding is adrenocortical hyperactivity. In this condition, blood Na is elevated while potassium is low. Potassium is elevated (wasted) in the urine. Observations at DDI indicate that Na and potassium levels in hair are commonly high in association with elevated levels of potentially toxic elements. The elevated Na and potassium levels are frequently concomitant with low levels of calcium and magnesium in hair. This apparent phenomenon requires further investigation.

Appropriate tests for Na status as an electrolyte are measurements of Na in whole blood and urine, and measurements of adrenocortical function.

Copper Normal

Hair Copper (Cu) levels are usually indicative of body status, except that exogenous contamination may occur giving a false normal (or false high). Common sources of contamination include: permanent solutions, dyes, bleaches, and swimming pools/hot tubs in which Cu compounds have been used as algacides.

Cu is an essential element that activates specific enzymes. Erythrocyte superoxide dismutase (SOD) is a Cu (and zinc) dependent enzyme; lysyl oxidase which catalyzes crosslinking of collagen is another Cu dependent enzyme. Adrenal catecholamine synthesis is Cu dependent, because the enzyme dopamine beta-hydroxylase, which catalyzes formation of norepinephrine from dopamine, requires Cu.

If hair Cu is in the normal range, this usually means tissue levels are in the normal range. However, under circumstances of contamination, a real Cu deficit could appear as a (false) normal. If symptoms of Cu deficiency are present, a whole blood or red blood cell elements analysis can be performed for confirmation of Cu status.

Zinc High

A result of high hair Zinc (Zn) may be indicative of low Zn in cells, and functional Zn deficiency. Zn can be displaced from proteins such as intracellular metallothionein by other metals, particularly cadmium, lead, copper, and mercury (Toxicology of Metals, 1994), resulting in paradoxically elevated hair Zn. Zn may also be high in hair as a result of the use of Zn-containing anti-dandruff shampoo. Rough or dry, flaky skin is a symptom of Zn deficiency, so it is not uncommon for Zn deficient patients to use an anti-dandruff shampoo. A result of high hair Zn warrants further testing to assess Zn status.

Zn is an essential element that is required in many very important biological processes. However, Zn can be toxic if exposure is excessive. Although very uncommon, high hair Zn might be indicative of Zn overload which could result from Zn contaminated water (galvanized pipes), welding or gross, chronic over-supplementation (100 mg/day). Other sources of Zn include: manufacture of brass, bronze, white paint, and pesticide production. Symptoms of Zn excess include: gastrointestinal disorders, decreased heme synthesis (copper deficiency), tachycardia, blurred vision, and hypothermia.

Confirmatory tests for Zn status are whole blood or packed red blood cell elements analysis, urine amino acid analysis, and serum ceruloplasmin (low with Zn induced copper deficiency).

Manganese Low

Hair Manganese (Mn) levels correlate well with Mn levels in other body tissues. Hair Mn levels are commonly low, in part due to low dietary Mn intake and the interaction of Mn with phosphates in the gut. Intestinal malabsorption also limits Mn uptake.

Mn is an essential element that is involved in energy metabolism, and bone and cartilage formation. Mn is an activator of many important enzymes including: mitochondrial superoxide dismutase, arginase, and pyruvate carboxylase.

Symptoms associated with Mn deficiency include: fatigue, lack of physical endurance, slow growth of fingernails and hair, impaired metabolism of bone and cartilage, dermatitis, weight loss, and reduced fertility. Increased allergic sensitivities and inflammation are often associated with low Mn. Seizures are occasionally reported to be associated with severe Mn deficiency.

An appropriate laboratory test to confirm Mn deficiency is whole blood elements analysis.

Cobalt Low

One can not determine vitamin B-12 status by use of hair analysis, and the clinical significance of low hair Cobalt (Co) levels is not known. Hair is analyzed for Co primarily for detection of excessive intake of the potentially toxic element.

There is little evidence that Co has an essential function in humans other than as an obligatory constituent of the vitamin B-12 molecule. Humans absorb Co as inorganic Co and as vitamin B-12; the body pools of each fluctuate independently. Humans cannot convert inorganic Co to vitamin B-12.

The dietary content of Co is highly variable, depending upon types of foods eaten, geographical location and type of soil. Vegetarians often have lower Co levels than meat eaters.

Appropriate tests for determination of vitamin B-12 status are the measurement of urine levels of methylmalonic acid (elevated with vitamin B-12 coenzyme deficiency/dysfunction), a quantitative blood assay for vitamin B-12, a urine amino acids analysis (several metabolic steps require vitamin B-12), and diet analysis.

Boron Low

Boron (B) is normally found in hair, but the correlations among dietary B intake, and tissue and hair levels of B have yet to be established. Recent studies clearly indicate that B has an important role in normal bone metabolism/density and may be needed for normal membrane function. In post-menopausal women consuming a very low B diet, B supplementation significantly lowered urinary excretion of calcium and magnesium and increased serum levels of estrogen (Environ. Health Perspect.; 102 Suppl.7: 59-63, 1994). Further research is in process to determine the clinical significance of hair B levels.

Lithium Low

Lithium (Li) is normally found in hair at very low levels. Hair Li correlates with high dosage of Li carbonate in patients treated for Affective Disorders. However, the clinical significance of low hair

Li levels is not certain at this time. Thus, hair Li is measured primarily for research purposes. Anecdotally, clinical feedback to DDI consultants suggests that low level Li supplementation may have some beneficial effects in patients with behavioral/emotional disorders. Li occurs almost universally in water and in the diet; excess Li is rapidly excreted in urine.

Li at low levels may have essential functions in humans. Intracellularly, Li inhibits the conversion of phosphorylated inositol to free inositol. In the nervous system this moderates neuronal excitability. Li also influences monamine neurotransmitter concentrations at the synapse (this function is increased when Li is used therapeutically for mania or bipolar illness).

A confirmatory test for low Li is measurement of Li in blood serum/plasma.

Selenium Low

Selenium (Se) is normally found in hair at very low levels, and several studies provide evidence that low hair Se is reflective of dietary intake and associated with cardiovascular disorders. Utilization of hair Se levels to assess nutritional status, however, is complicated by the fact that use of Se- or sulfur-containing shampoo markedly increases hair Se (externally) and can give a false high value.

Se is an extremely important essential element due to its antioxidative function as an obligatory component of the enzyme glutathione peroxidase. Se is also protective in its capacity to bind and "inactivate" mercury, and Se is an essential cofactor in the deiodination of T-4 to active T-3 (thyroid hormone). Some conditions of functional hypothyroidism therefore may be due to Se deficiency (Nature; 349:438-440, 1991); this is of particular concern with mercury exposure. Studies have also indicated significant inverse correlations between Se and heart disease, cancer, and asthma.

Selenium deficiency is common and can result from low dietary intake of Se or vitamin E, and exposure to toxic metals, pesticides/herbicides and chemical solvents.

Symptoms of Se deficiency are similar to that of vitamin E deficiency and include muscle aches, increased inflammatory response, loss of body weight, alopecia, listlessness, skeletal and muscular degeneration, growth stunting, and depressed immune function.

Confirmatory tests for Se deficiency are Se content of packed red blood cells, and activity of glutathione peroxidase in red blood cells.

Total Toxic Element Indication

The potentially toxic elements vary considerably with respect to their relative toxicities. The accumulation of more than one of the most toxic elements may have synergistic adverse effects, even if the level of each individual element is not strikingly high. Therefore, we present a total toxic element "score" which is estimated using a weighted average based upon relative toxicity. For example, the combined presence of lead and mercury will give a higher total score than that of the combination of silver and beryllium.

Lab number: **H061205-0496-1**
Patient: **Katherine Spehar**

Hair

Page: 6
Client: **24237**
